arXiv:1611.09251v1 [math.DS] 28 Nov 2016

Monotone Dynamical Systems with Polyhedral Order Cones and Dense Periodic Points

Morris W. Hirsch
Department of Mathematics
University of Wisconsin at Madison
University of California at Berkeley

November 29, 2016

Abstract

Let $X \subset \mathbb{R}^n$ be a nonempty whose interior is connected and dense in X. Assume $T\colon X \approx X$ is injective, continuous, and monotone for the order defined by a closed convex cone $K \subset \mathbb{R}^n$.

Theorem. If K is polyhedral and the set of periodic points is dense, then T is periodic.

Contents

1 Introduction **1**
Notation 2

2 Proof of the Main Theorem **3**

1 Introduction

$K \subset \mathbb{R}^n$ (Euclidean n-space) is a closed convex cone that is *solid* (has nonempty interior $\mathsf{int}\,(K)$), and *pointed* (contains no affine line). Briefly: K is a *solid order cone* in $\mathbb{R}^n$.

$\mathbb{R}^n$ is given the (partial) order $\geq$ determined by K, referred to as the *K-order*:

$$y \geq x \iff y - x \in K,$$

$X \subset \mathbb{R}^n$ is a nonempty set whose interior $\mathsf{int}\,(X)$ is connected and dense in X.
$T\colon X \to X$ is an injective continuous map that is *monotone* for the K-order:

$$x \geq y \implies Tx \geq Ty.$$

x has *period* k if k is a positive integer and $T^k x = x$. The set of these points is denoted by $\mathcal{P}_k = \mathcal{P}_k(T)$, and the set of periodic pointsby $\mathcal{P} = \mathcal{P}(T) = \bigcup_k \mathcal{P}_k$.

T is *periodic* if $X = \mathcal{P}_k$, and *pointwise periodic* if $X = \mathcal{P}$.

Our main concern is the following speculation:

Conjecture. *If $\mathcal{P}$ is dense in X, then T is periodic.*

We prove this for K a *polyhedron*: the intersection of finitely many closed affine halfspaces of $\mathbb{R}^n$.

Theorem 1 (Main)**.** *If K is a polyhedron and $\mathcal{P}$ is dense in X, then T is periodic.*

The following result of D. Montgomery [3][1] is crucial to the proof:

Theorem 2 (Montgomery)**.** *Every pointwise periodic homeomorphism of a connected manifold is periodic.*

This implies a sharper version of Theorem 1 for analytic maps:

Theorem 3. *If K is a polyhedron and T is analytic but not periodic, $\mathcal{P}$ is nowhere dense.*

Two observations simplify Theorem 1:

- The hypothesis holds if $\mathcal{P}$ is dense in $\mathsf{int}\,(X)$, because $\mathsf{int}\,(X)$ is dense in X.
- The conclusion holds if $\mathsf{int}\,(X) \subset \mathcal{P}$.

  For then Montgomery's Theorem implies $\mathsf{int}\,(X) \subset \mathcal{P}_k$, hence $X \subset \mathcal{P}_k$ because $\mathsf{int}\,(X)$ is dense in X and $\mathcal{P}_k$ is closed in X,

Notation

i, j, k, l always denote positive integers, and $a, b, p, q, u, v, x, y, z$ points of $\mathbb{R}^n$.

$x \preceq y$ is a synonym for $y \succeq x$. If $x \preceq y$ and $x \neq y$ we write $x \prec$ or $y \succ x$.

The relations $x \ll y$ and $y \gg x$ mean $y - x \in \mathsf{int}\,(K)$.

A set S is *totally ordered* if $x, y \in S \implies x \preceq y$ or $x \succeq y$.

If $x \preceq y$, the *order interval* $[x, y]$ is $\{z\colon\ x \preceq z \preceq y\} = K_x \cap -K_y$.

The translation of K by $x \in \mathbb{R}^n$ is $K_x := \{w + x, w \in K.\}$

The image of a set or point ξ under a map H is denoted by $H\xi$ or $H(\xi)$. A set S is *positively invariant* under H if $HS \subset S$, *invariant* if $H\xi = \xi$, and *periodically invariant* if $H^k\xi = \xi$.

[1] See also S. Kaul [2]

2 Proof of the Main Theorem

We derive three topological consequences, valid even if K is not polyhedral, from the standing assumptions that T is monotone and $\mathcal{P}$ is dense:

Proposition 4. *Assune $p, q \in \mathcal{P}_k$ are such that*

$$p \ll q, \qquad p, q \in \mathcal{P}_k. \qquad [p, q] \subset X.$$

Then $T^k([p, q] = [p, q]$.

Proof. It suffices to take $k = 1$. Evidently $T\mathcal{P} = \mathcal{P}$, and $T[p, q] \subset [p, q]$ because T is monotone, whence $\mathsf{int}\,([p, q]) \cap \mathcal{P}$ is positively invariant under T. The conclusion follows because $\mathsf{int}\,[p, q] \cap \mathcal{P}$ is dense in $[p, q]$ and T is continuous. ∎

Proposition 5. *Assume $u, v \in \mathcal{P}_k, u \ll v$, and $[u, v] \subset X$. There is a compact arc $J \subset \mathcal{P}_k \cap [u, v]$ joining u to v that is totally ordered by $\ll$.*[2]

Proof. An application of Zorn's Lemma yields a maximal set $J \subset [u, v] \cap P$ such that: J is totally ordered by $\ll$, $u = \max J$, $v = \min J$. Maximality implies J is compact and connected and $u, v \in J$, so J is an arc (Wilder [5], Theorem I.11.23). ∎

Proposition 6. *Let $M \subset X$ be a topological manifold of dimension $n - 1$ without boundary, closed in X.*

(i) *$\mathcal{P}$ is dense in M.*

(ii) *If M is periodically invariant, it has a neighborhood base $\mathcal{W}$ of periodically invariant open subsets.*

(iii) *$\mathcal{P}$ is dense in each $W \in \mathcal{W}$.*

Proof. Lefschetz duality (Spanier [4]) shows that every point of M has arbitrarily small open ball neighborhoods $W \subset X$ separated by M into two disjoint open sets U and V:

$$W \setminus M = U \cup V, \quad U \cap V = \varnothing. \tag{1}$$

Since $\mathcal{P}$ is dense in W there exist $u, v \in \mathcal{P} \cap U, v \in \mathcal{P} \cap V$ such that

$$u \ll v, \qquad [u, v] \subset W.$$

There is an arc $J \subset \mathcal{P} \cap [u, v]$ joining u and v (Proposition 5). Because J is separated by M, there exists $p \in J \cap M$, implying $p \in W \cap \mathcal{P}$. This proves (i), assertion (ii) follows because $M \cap \mathsf{int}\,([u, v])$ is periodically invariant, and (iii) holds because $\mathcal{P}$ is dense M and W is open in M. ∎

Let $\mathcal{T}(m)$ stand for the statement of Theorem 1 for the case $n = m$. Then $\mathcal{T}(0)$ is trivial, and we use the following inductive hypothesis:

[2] This result is adapted from Hirsch & Smith [1], Theorems 5,11 & 5,15.

Hypothesis 7 (INDUCTION)**.** *$n \geq 1$ and $\mathcal{T}(n-1)$ holds.*

Let $Q \subset \mathbb{R}^n$ be a compact n-dimensional polyhedral cell, such as $[p,q]$ with $p \ll q$. Its boundary ∂Q is the union of finitely many convex compact $(n-1)$-cells, the *faces* of Q. Each face F is the intersection of $\partial[p,q]$ with a unique affine hyperplane E^{n-1}. The corresponding *open face* is $F^\circ := F \setminus \partial F$, an open $(n-1)$-cell in E^{n-1}. Distinct open faces are disjoint, and their union is dense and open in ∂Q.

Proposition 8. *Assume $p, q \in \mathcal{P}_k,\ p \ll q,\ [p,q] \subset X$. Then $T|\partial[p,q]$ is periodic.*

Proof. The $(n-1)$-dimensional compact polyhedron $\partial[p,q]$ is invariant under T^k (Proposition 4). Each open face $F^\circ \subset \partial[p,q]$ is a union of periodically invariant open subsets W_λ, each an $(n-1)$-manifold without boundary, with $\mathcal{P}$ dense in W_λ (Proposition 6).

The maps $T|W_\lambda$ are periodic by the Induction Hypothesis, so $F^\circ \subset \mathcal{P}$. Montgomery's Theorem implies $T|F^\circ$ is periodic, so $T|F$ is also periodic. Since $\partial[p,q]$ is the union of the finitely many faces F, it follows that $T|\partial[p,q]$ is periodic. ∎

To complete the inductive proof of the Main Theorem, it suffices by Montgomery's theorem to prove that an arbitrary $x \in X$ is periodic. As X is open in $\mathbb{R}^n$ and $\mathcal{P}$ is dense in X, there is an order interval $[a,b] \subset X$ such that

$$a \ll x \ll b, \qquad a, b \in \mathcal{P}_k.$$

By Proposition 5, a and b are the endpoints of a compact arc $J \subset \mathcal{P}_k \cap [a,b]$, totally ordered by $\ll$. Define $p, q \in J$:

$$p := \sup\{y \in J :\ y \preceq x\}, \qquad q := \inf\{y \in J :\ y \succeq x\}.$$

If $p = q = x$ then $x \in \mathcal{P}_k$. Otherwise $p \ll q$ and $x \in \partial[p,q]$, whence $x \in \mathcal{P}$ by Proposition 8. ∎

References

[1] M. Hirsch & H. Smith, *Monotone Dynamical Systems,* "Handbook of Differential Equations," volume 2, chapter 4. A. Cañada, P. Drabek & A. Fonda editors. Elsevier North Holland 2005.

[2] S. Kaul, *On pointwise periodic transformation groups,* Proc. Amer. Math. Soc. **27** (1971), 391-394.

[3] D. Montgomery, *Pointwise periodic homeomorphisms,* American Journal of Mathematics **59** (1937), 118-120.

[4] E. Spanier, "Algebraic Topology," McGraw Hill 1966.

[5] R. Wilder, "Topology of Manifolds," Amer. Math. Soc. 1949.